arXiv:1304.0546v1 [math.MG] 2 Apr 2013

Volumes and geodesic ball packings to the regular prism tilings in $\widetilde{\mathbf{SL_2R}}$ space [*]

Dedicated to Prof. Lajos Tamássy on His 90^{th} Birthday

Emil Molnár and Jenő Szirmai
Budapest University of Technology and
Economics Institute of Mathematics,
Department of Geometry
Budapest, P. O. Box: 91, H-1521
emolnar@math.bme.hu, szirmai@math.bme.hu

April 3, 2013

Abstract

After having investigated the regular prisms and prism tilings in $\widetilde{\mathbf{SL_2R}}$ space in previous work [13] of the second author, we consider the problem of geodesic ball packings related to those tilings and their symmetry groups $\mathbf{pq2_1}$. $\widetilde{\mathbf{SL_2R}}$ geometry is one among the eight Thurston geometries that can be derived from the 3-dimensional Lie group of all 2×2 real matrices with determinant one.

In this paper we consider geodesic spheres and balls in $\widetilde{\mathbf{SL_2R}}$ space (even in $\mathbf{SL_2R}$), if its radius $\rho \in [0, \frac{\pi}{2})$ and determine their volumes. Moreover, we consider the prisms of the above space and compute their volumes, define the notion of the geodesic ball packing and its density. We develop a procedure to determine the densities of the densest geodesic ball packings for the regular prism tilings, moreover for their group $\mathbf{pq2_1}$, and apply this

---

[*] Mathematics Subject Classification 2010: 52C17, 52C22, 52B15, 53A35, 51M20.
Key words and phrases: Thurston geometries, $\widetilde{\mathbf{SL_2R}}$ geometry, ball packings, tiling, regular prism, volume in $\widetilde{\mathbf{SL_2R}}$ space.

algorithm to them. Surprisingly we found a very dense packing in $\widetilde{\mathbf{SL_2R}}$ (Theorem 4.1), denser than those in the hyperbolic space $\mathbf{H}^3$.

We use for the computations and visualization of $\widetilde{\mathbf{SL_2R}}$ space its projective model introduced by the first author in [4].

1 On $\widetilde{\mathrm{SL_2R}}$ geometry

The real 2×2 matrices $\begin{pmatrix} d & b \\ c & a \end{pmatrix}$ with unit determinant $ad - bc = 1$ constitute a Lie transformation group by the usual product operation, taken to act on row matrices as on point coordinates on the right as follows

$$(z^0, z^1) \begin{pmatrix} d & b \\ c & a \end{pmatrix} = (z^0 d + z^1 c, z^0 b + z^1 a) = (w^0, w^1). \tag{1.1}$$

This group is a 3-dimensional manifold, because of its 3 independent real coordinates and with its usual neighbourhood topology [8], [15]. In order to model the above structure in the projective sphere $\mathcal{PS}^3$ and in projective space $\mathcal{P}^3$ (see [4]) we introduce the new projective coordinates (x^0, x^1, x^2, x^3) where

$$a := x^0 + x^3, \ b := x^1 + x^2, \ c := -x^1 + x^2, \ d := x^0 - x^3,$$

with positive, then the non-zero multiplicative equivalence as a projective freedom in $\mathcal{PS}^3$ and in $\mathcal{P}^3$, respectively. Meanwhile we turn to the proportionality $\mathbf{SL_2R} \sim \mathbf{PSL_2R}$, natural in this context. Then it follows that

$$0 > bc - ad = -x^0x^0 - x^1x^1 + x^2x^2 + x^3x^3 \tag{1.2}$$

describes the interior of the above one-sheeted hyperboloid solid $\mathcal{H}$ in the usual Euclidean coordinate simplex, with the origin $E_0(1;0;0;0)$ and the ideal points of the axes $E_1^\infty(0;1;0;0)$, $E_2^\infty(0;0;1;0)$, $E_3^\infty(0;0;0;1)$. We consider the collineation group $\mathbf{G}_*$ that acts on the projective space $\mathcal{P}^3$ and preserves a polarity, i.e. a scalar product of signature $(- - ++)$, this group leaves the one sheeted hyperboloid solid $\mathcal{H}$ invariant. We have to choose an appropriate subgroup $\mathbf{G}$ of $\mathbf{G}_*$ as isometry group, then the universal covering group and space $\widetilde{\mathcal{H}}$ of $\mathcal{H}$ will be the hyperboloid model of $\widetilde{\mathbf{SL_2R}}$ (see [4]).

The specific isometries $\mathbf{S}(\phi)$ $(\phi \in \mathbf{R})$ constitute a one parameter group given by the matrices

$$\mathbf{S}(\phi): \ (s_i^j(\phi)) = \begin{pmatrix} \cos\phi & \sin\phi & 0 & 0 \\ -\sin\phi & \cos\phi & 0 & 0 \\ 0 & 0 & \cos\phi & -\sin\phi \\ 0 & 0 & \sin\phi & \cos\phi \end{pmatrix}. \tag{1.3}$$

The elements of $\mathbf{S}(\phi)$ are the so-called *fibre translations*. We obtain a unique fibre line to each $X(x^0; x^1; x^2; x^3) \in \widetilde{\mathcal{H}}$ as the orbit by right action of $\mathbf{S}(\phi)$ on X. The coordinates of points lying on the fibre line through X can be expressed as the images of X by $\mathbf{S}(\phi)$:

$$\begin{gathered}(x^0; x^1; x^2; x^3) \xrightarrow{\mathbf{S}(\phi)} (x^0\cos\phi - x^1\sin\phi; x^0\sin\phi + x^1\cos\phi; \\ x^2\cos\phi + x^3\sin\phi; -x^2\sin\phi + x^3\cos\phi).\end{gathered} \tag{1.4}$$

The points of a fibre line through X by usual inhomogeneous Euclidean coordinates $x = \frac{x^1}{x^0}$, $y = \frac{x^2}{x^0}$, $z = \frac{x^3}{x^0}$, $x^0 \neq 0$ are given by

$$(1; x; y; z) \xrightarrow{\mathbf{S}(\phi)} \Big(1; \frac{x + \tan\phi}{1 - x\tan\phi}; \frac{y + z\tan\phi}{1 - x\tan\phi}; \frac{z - y\tan\phi}{1 - x\tan\phi}\Big) \tag{1.5}$$

for the projective space $\mathcal{P}^3$, where ideal points (at infinity) conventionally occur. The π periodicity of the above maps can be seen from the formula (1.5) e.g. if $-\frac{\pi}{2} < \phi < \frac{\pi}{2}$ then $-\infty < x < \infty$. In (1.3) and (1.4) we can see the 2π periodicity of ϕ, moreover the (logical) extension to $\phi \in \mathbf{R}$, as real parameter, to have the



Figure 1: The hyperboloid model

universal covers $\widetilde{\mathcal{H}}$ and $\widetilde{\mathbf{SL_2R}}$, respectively, through the projective sphere $\mathcal{PS}^3$. The elements of the isometry group of $\mathbf{SL_2R}$ (and so by the above extension the isometries of $\widetilde{\mathbf{SL_2R}}$) can be described by the matrix (a_i^j) (see [4] and [5])

$$(a_i^j) = \begin{pmatrix} a_0^0 & a_0^1 & a_0^2 & a_0^3 \\ \mp a_0^1 & \pm a_0^0 & \pm a_0^3 & \mp a_0^2 \\ a_2^0 & a_2^1 & a_2^2 & a_2^3 \\ \pm a_2^1 & \mp a_2^0 & \mp a_2^3 & \pm a_2^2 \end{pmatrix} \quad \text{where}$$

$$-(a_0^0)^2 - (a_0^1)^2 + (a_0^2)^2 + (a_0^3)^2 = -1, \;\; -(a_2^0)^2 - (a_2^1)^2 + (a_2^2)^2 + (a_2^3)^2 = 1,$$
$$-a_0^0 a_2^0 - a_0^1 a_2^1 + a_0^2 a_2^2 + a_0^3 a_2^3 = 0 = -a_0^0 a_2^1 + a_0^1 a_2^0 - a_0^2 a_2^3 + a_0^3 a_2^2. \tag{1.6}$$

We define the *translation group* $\mathbf{G}_T$, as a subgroup of the isometry group of $\mathbf{SL_2R}$, the isometries acting transitively on the points of $\mathcal{H}$ and by the above extension on the points of $\widetilde{\mathcal{H}}$. $\mathbf{G}_T$ maps the origin $E_0(1;0;0;0)$ onto $X(x^0;x^1;x^2;x^3)$. These isometries and their inverses (up to a positive determinant factor) can be given by the following (t_i^j) and $T_j^k = (t_i^j)^{-1}$ matrices:

$$\begin{gathered}\mathbf{T}: \; (t_i^j) = \begin{pmatrix} x^0 & x^1 & x^2 & x^3 \\ -x^1 & x^0 & x^3 & -x^2 \\ x^2 & x^3 & x^0 & x^1 \\ x^3 & -x^2 & -x^1 & x^0 \end{pmatrix}, \\ \mathbf{T}^{-1}: \; (T_j^k) = \begin{pmatrix} x^0 & -x^1 & -x^2 & -x^3 \\ x^1 & x^0 & -x^3 & x^2 \\ -x^2 & -x^3 & x^0 & -x^1 \\ -x^3 & x^2 & x^1 & x^0 \end{pmatrix}.\end{gathered} \tag{1.7}$$

The rotation about the fibre line through the origin $E_0(1;0;0;0)$ by angle ω $(-\pi < \omega \leq \pi)$ can be expressed by the following matrix (see [4])

$$\mathbf{R}_{E_O}(\omega): \; (r_i^j(E_0,\omega)) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \cos\omega & \sin\omega \\ 0 & 0 & -\sin\omega & \cos\omega \end{pmatrix}, \tag{1.8}$$

and the rotation $\mathbf{R}_X(\omega)$ with matrix : $(r_i^j(X,\omega))$ about the fibre line through $X(x^0;x^1;x^2;x^3)$ by angle ω can be derived by formulas (1.7) and (1.8) where $\mathbf{R}_X(\omega) = \mathbf{T}^{-1}\mathbf{R}_{E_O}(\omega)\mathbf{T}$. Thus the above rotation $\mathbf{R}_X(\omega)$, with a specific X

$(\cosh r, 0, \sinh r, 0) \sim (1, 0, \tanh r, 0)$ has the important matrix

$$\begin{pmatrix}
\begin{array}{c} 1+\sinh^2 r - \\ - \sinh^2 r \cos\omega \end{array} & \sinh^2 r \sin\omega & \begin{array}{c} \dfrac{1}{2}\sinh 2r - \\ -\dfrac{1}{2}\sinh 2r \cos\omega \end{array} & -\frac{1}{2}\sinh 2r \sin\omega \\
- \sinh^2 r \sin\omega & \begin{array}{c} 1+\sinh^2 r - \\ - \sinh^2 r \cos\omega \end{array} & -\frac{1}{2}\sinh 2r \sin\omega & \begin{array}{c} -\dfrac{1}{2}\sinh 2r + \\ +\dfrac{1}{2}\sinh 2r \cos\omega \end{array} \\
\begin{array}{c} -\dfrac{1}{2}\sinh 2r + \\ +\dfrac{1}{2}\sinh 2r \cos\omega \end{array} & -\frac{1}{2}\sinh 2r \sin\omega & \begin{array}{c} 1-\cosh^2 r + \\ + \cosh^2 r \cos\omega \end{array} & \cosh^2 r \sin\omega \\
-\frac{1}{2}\sinh 2r \sin\omega & \begin{array}{c} \dfrac{1}{2}\sinh 2r - \\ -\dfrac{1}{2}\sinh 2r \cos\omega \end{array} & - \cosh^2 r \sin\omega & \begin{array}{c} 1-\cosh^2 r + \\ + \cosh^2 r \cos\omega \end{array}
\end{pmatrix} \tag{1.8'}$$

Horizontal intersection of the hyperboloid solid $\mathcal{H}$ with the plane $E_0 E_2^\infty E_3^\infty$ provides the *base plane* of the model $\widetilde{\mathcal{H}} = \widetilde{\mathbf{SL_2R}}$. The fibre through X intersects the base plane $z^1 = x = 0$ in a foot point

$$Z(z^0 = x^0x^0 + x^1x^1; z^1 = 0; z^2 = x^0x^2 - x^1x^3; z^3 = x^0x^3 + x^1x^2). \tag{1.9}$$

We generally introduce a so-called hyperboloid parametrization by [4] as follows

$$\begin{aligned}
x^0 &= \cosh r \cos\phi, \\
x^1 &= \cosh r \sin\phi, \\
x^2 &= \sinh r \cos(\theta - \phi), \\
x^3 &= \sinh r \sin(\theta - \phi),
\end{aligned} \tag{1.10}$$

where (r, θ) are the polar coordinates of the base plane and ϕ is just the fibre coordinate. We note that

$$-x^0x^0 - x^1x^1 + x^2x^2 + x^3x^3 = -\cosh^2 r + \sinh^2 r = -1 < 0.$$

The inhomogeneous coordinates in (1.11), that will play an important role in the later $\mathbf{E}^3$-visualization of the prism tilings in $\widetilde{\mathbf{SL_2R}}$, are given by

$$\begin{aligned} x &= \frac{x^1}{x^0} = \tan\phi, \\ y &= \frac{x^2}{x^0} = \tanh r \frac{\cos(\theta-\phi)}{\cos\phi}, \\ z &= \frac{x^3}{x^0} = \tanh r \frac{\sin(\theta-\phi)}{\cos\phi}. \end{aligned} \tag{1.11}$$

The infinitesimal arc-length square can be derived by the standard method, called pull back into the origin. By acting of (1.7) on the differentials $(dx^0; dx^1; dx^2; dx^3)$, we obtain by [4], [2] and [5] that in this parametrization the infinitesimal arc-length-square at any point of $\widetilde{\mathbf{SL_2R}}$ is the following:

$$(ds)^2 = (dr)^2 + \cosh^2 r \sinh^2 r (d\theta)^2 + \big[(d\phi) + \sinh^2 r (d\theta)\big]^2. \tag{1.12}$$

Hence we get the symmetric metric tensor field g_{ij} on $\widetilde{\mathbf{SL_2R}}$ by components:

$$g_{ij} := \begin{pmatrix} 1 & 0 & 0 \\ 0 & \sinh^2 r(\sinh^2 r + \cosh^2 r) & \sinh^2 r \\ 0 & \sinh^2 r & 1 \end{pmatrix}, \tag{1.13}$$

and

$$dV = \sqrt{\det(g_{ij})}\, dr\, d\theta\, d\phi = \frac{1}{2}\sinh(2r) dr\, d\theta\, d\phi$$

as the volume element in hyperboloid coordinates. The geodesic curves of $\widetilde{\mathbf{SL_2R}}$ are generally defined as having locally minimal arc length between any two of their (near enough) points.

By (1.13) the second order differential equation system of the $\widetilde{\mathbf{SL_2R}}$ geodesic curve is the following:

$$\begin{aligned} \ddot{r} &= \sinh(2r)\dot{\theta}\dot{\phi} + \frac{1}{2}\big(\sinh(4r) - \sinh(2r)\big)\dot{\theta}\dot{\theta}, \\ \ddot{\phi} &= 2\dot{r}\tanh(r)(2\sinh^2(r)\dot{\theta} + \dot{\phi}), \\ \ddot{\theta} &= \frac{2\dot{r}}{\sinh(2r)}\big((3\cosh(2r) - 1)\dot{\theta} + 2\dot{\phi}\big). \end{aligned} \tag{1.14}$$

We can assume, that the starting point of a geodesic curve is the origin $(1, 0, 0, 0)$, because we can transform a curve into an arbitrary starting point. Moreover, $r(0) = 0,\ \phi(0) = 0,\ \theta(0) = 0,\ \dot{r}(0) = \cos(\alpha),\ \dot{\phi}(0) = \sin(\alpha) = -\dot{\theta}(0)$ in Table 1 for the solution of (1.14) and so the unit velocity will be achieved.

Table 1

Types	
$0 \leq \alpha < \frac{\pi}{4}$ $(\mathbf{H}^2 - \text{like direction})$	$r(s,\alpha) = \operatorname{arsinh}\Big(\frac{\cos\alpha}{\sqrt{\cos 2\alpha}}\sinh(s\sqrt{\cos 2\alpha})\Big)$ $\theta(s,\alpha) = -\arctan\Big(\frac{\sin\alpha}{\sqrt{\cos 2\alpha}}\tanh(s\sqrt{\cos 2\alpha})\Big)$ $\phi(s,\alpha) = 2\sin\alpha s + \theta(s,\alpha)$
$\alpha = \frac{\pi}{4}$ (light direction)	$r(s,\alpha) = \operatorname{arsinh}\Big(\frac{\sqrt{2}}{2}s\Big)$ $\theta(s,\alpha) = -\arctan\Big(\frac{\sqrt{2}}{2}s\Big)$ $\phi(s,\alpha) = \sqrt{2}s + \theta(s,\alpha)$
$\frac{\pi}{4} < \alpha \leq \frac{\pi}{2}$ $(\text{fibre} - \text{like direction})$	$r(s,\alpha) = \operatorname{arsinh}\Big(\frac{\cos\alpha}{\sqrt{-\cos 2\alpha}}\sin(s\sqrt{-\cos 2\alpha})\Big)$ $\theta(s,\alpha) = -\arctan\Big(\frac{\sin\alpha}{\sqrt{-\cos 2\alpha}}\tan(s\sqrt{-\cos 2\alpha})\Big)$ $\phi(s,\alpha) = 2\sin\alpha s + \theta(s,\alpha)$

The equation of the geodesic curve in the hyperboloid model has been determined in [2], with the usual geographical sphere coordiantes (λ,α), as longitude and altitude, respectively, from the general starting position of (1.10), (1.11), $(-\pi < \lambda \leq \pi,\ -\frac{\pi}{2} \leq \alpha \leq \frac{\pi}{2})$, and the arc-length parameter $0 \leq s \in \mathbf{R}$. The Euclidean coordinates $X(s,\lambda,\alpha)$, $Y(s,\lambda,\alpha)$, $Z(s,\lambda,\alpha)$ of the geodesic curves can be determined by substituting the results of Table 1 (see [2]) into the equations (1.10) and (1.11) as follows :

$$\begin{gathered} X(s,\lambda,\alpha) = \tan\left(\phi(s,\alpha)\right), \\ Y(s,\lambda,\alpha) = \frac{\tanh\left(r(s,\alpha)\right)}{\cos\left(\phi(s,\alpha)\right)}\cos\big[\theta(s,\alpha) - \phi(s,\alpha) + \lambda\big], \end{gathered} \tag{1.15}$$

$$Z(s,\lambda,\alpha) = \frac{\tanh\left(r(s,\alpha)\right)}{\sin\left(\phi(s,\alpha)\right)}\cos\big[\theta(s,\alpha) - \phi(s,\alpha) + \lambda\big].$$

2 Geodesic balls in $\widetilde{\mathbf{SL_2R}}$ space

Definition 2.1 *The distance $d(P_1, P_2)$ between the points P_1 and P_2 is defined by the arc length of the geodesic curve from P_1 to P_2.*

The numerical approximation of the distance $d(O, P)$, by Table 1 and (1.15) for given $P(X, Y, Z)$ from the origin O, will not be detailed here.

Definition 2.2 *The geodesic sphere of radius ρ (denoted by $S_{P_1}(\rho)$) with the center at point P_1 is defined as the set of all points P_2 in space with the condition $d(P_1, P_2) = \rho$. Moreover, we require that the geodesic sphere is a simply connected surface without selfintersection in $\widetilde{\mathbf{SL_2R}}$.*

Definition 2.3 *The body of the geodesic sphere of centre P_1 and of radius ρ in the $\widetilde{\mathbf{SL_2R}}$ space is called geodesic ball, denoted by $B_{P_1}(\rho)$, i.e., $Q \in B_{P_1}(\rho)$ iff $0 \leq d(P_1, Q) \leq \rho$.*

Remark 2.1 *We choose $O(1, 0, 0, 0)$ as centre of the sphere and its ball by the homogeneity of $\widetilde{\mathbf{SL_2R}}$.*

Fig. 2.a shows a geodesic sphere of radius $\rho = 1.3$ with centre O and Fig. 2.b shows its intersection with the (x, z) plane. From (1.15) it follows that $S(\rho)$ is a simply connected surface in $\mathbf{E}^3$ and $\widetilde{\mathbf{SL_2R}}$, respectively, if $\rho \in [0, \frac{\pi}{2})$. If$\rho \geq \frac{\pi}{2}$ then the universal cover should be discussed. *Therefore, we consider the geodesic spheres and balls of radius $\rho \in [0, \frac{\pi}{2})$.* These will be satisfactory for our prism tilings.

2.1 The volume of a geodesic ball

The volume formula of the geodesic ball $B(\rho)$ of radius ρ follows from the metric tensor g_{ij}. We obtain the connection between the hyperboloid coordinates (r, θ, ϕ) and the geographical coordinates (s, λ, α) by the Table 1 and by the equation (1.15). Therefore, the volume of the geodesic ball of radius ρ can be computed by the following integral:



Figure 2: a, b

Theorem 2.1

$$
\begin{gathered}
Vol(B(\rho)) = \int_B \frac{1}{2}\sinh(2r)\; dr\; d\theta\; d\phi = \\
= 4\pi \int_0^{\rho} \int_0^{\frac{\pi}{4}} \frac{1}{2}\sinh(2r(s,\alpha))|J_1|\; d\alpha\; ds + \\
+4\pi \int_0^{\rho} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{1}{2}\sinh(2r(s,\alpha))|J_2|\; d\alpha\; ds
\end{gathered}
\tag{2.1}
$$

where $|J_1| = \begin{vmatrix} \frac{\partial r}{\partial s} & \frac{\partial r}{\partial \alpha} \\ \frac{\partial \phi}{\partial s} & \frac{\partial \phi}{\partial \alpha} \end{vmatrix}$ *and similarly* $|J_2|$ *(by Table 1 and* $\frac{\partial \theta}{\partial \lambda} = 1$*) are the corresponding Jacobians.*

The complicated formulas above need numerical approximation by computer. In Table 2 we have collected the volumes of the geodesic balls for some radii (Fig. 3).



Figure 3: The increasing function $\rho \mapsto Vol(B(\rho))$.

Table 2	
ρ	$Vol(B(\rho))$
$\rho = 0.1$	≈ 0.004202769
$\rho = 0.5$	≈ 0.568541467
$\rho = 1$	≈ 5.761162741
$\rho = 1.3$	≈ 15.54409646
$\rho \to \frac{\pi}{2}$	≈ 34.17279403

3 Regular prisms and prism tilings in $\widetilde{\mathrm{SL}_2\mathrm{R}}$ space

In the paper [13] we have defined and described the regular prisms and prism tilings in $\widetilde{\mathbf{SL_2R}}$ space. These will be summarized in this section.

Definition 3.1 *Let $\mathcal{P}^i$ be an infinite solid that is bounded by certain surfaces in the model space of $\widetilde{\mathbf{SL_2R}}$ that will be determined (later on) by „side fibre lines” passing through the vertices of a regular p-gon ($\mathcal{P}^b$) lying in the „base plane”.*

The images of solids $\mathcal{P}^i$ by $\widetilde{\mathbf{SL_2R}}$ isometries are called infinite regular p-sided $\widetilde{\mathbf{SL_2R}}$ prisms. Here regular means that the side surfaces are congruent to each other under $\widetilde{\mathbf{SL_2R}}$ rotations about a fiber line (e.g. through the origin).

The common part of $\mathcal{P}^i$ with the base plane is the *base figure* of $\mathcal{P}^i$ that is denoted by $\mathcal{P}$ and its vertices coincide with the vertices of $\mathcal{P}^b$ **but $\mathcal{P}$ is not assumed to be a polygon**.

Definition 3.2 *A bounded regular p-sided prism in $\widetilde{\mathbf{SL_2R}}$ is analogously defined if the face of the base figure $\mathcal{P}$ and its translated copy $\mathcal{P}^t$ by a fibre translation by (1.3) and so (1.5) are also introduced. The faces $\mathcal{P}$ and $\mathcal{P}^t$ are called cover faces.*

Remark 3.1 *All cross-sections of a prism „parallel" (the intersecting planes are generated by $\widetilde{\mathbf{SL_2R}}$ fibre translations from the base plane) to the base faces are congruent. Prisms are named for their base, e.g. a prism with a pentagonal base is called a pentagonal prism.*

In [13] we have considered regular prism tilings $\mathcal{T}_p(q)$ by prisms $\mathcal{P}_p(q)$ where q pieces regularly meet at each side edge by q-rotation.

The following theorem have been proved:

Theorem 3.1 *There exist regular infinite prism tilings $\mathcal{T}_p^i(q)$ in $\widetilde{\mathbf{SL_2R}}$ for each $3 \leq p \in \mathbb{N}$ where $\frac{2p}{p-2} < q \in \mathbb{N}$. For bounded prisms, these are not face-to-face.*

In this paper we consider the above regular bounded prism tilings $\mathcal{T}_p(q)$. Then each vertex of a tiling is a proper point of $\widetilde{\mathbf{SL_2R}}$, and the prism is a „topological polyhedron" having at each vertex one „p-gonal cover face" (it is not a polygon at all) and two skew quadrangles which lie on certain side surfaces in the model. Let $\mathcal{P}_p(q)$ be one of the tiles where $\mathcal{P}$ (and so $\mathcal{P}^b$ as well) is centered at the origin with vertices $A_1A_2A_3\dots A_p$ in the base plane. It is clear that the side curves $c_{A_iA_{i+1}}$ $(i = 1\dots p,\ A_{p+1} \equiv A_1)$ of the base figure are derived from each other by $\frac{2\pi}{p}$ rotation about the vertical x axis, so there are congruent in $\widetilde{\mathbf{SL_2R}}$ sense. The corresponding vertices $B_1B_2B_3\dots B_p$ are generated by a fibre translation τ given by (1.3) with parameter $0 < \Phi \in \mathbb{R} \setminus \{0\}$. The fibre lines through the vertices A_iB_i are denoted by $f_i,\ (i = 1, \dots, p)$ and the fibre line through the "midpoint" H of the curve $c_{A_1A_p}$ is denoted by f_0. This f_0 will be a half-screw axis of $\mathbf{ab}$ as follows.



Figure 4:

The tiling $\mathcal{T}_p(q)$ is generated by its discrete isometry group $\Gamma_p(q) = \mathbf{pq2_1} \subset Isom(\widetilde{\mathbf{SL_2R}})$ which is given by its fundamental domain $A_1A_2OA_1^{\mathbf{s}}A_2^{\mathbf{s}}O^{\mathbf{s}}$ a „topological polyhedron" and the group presentation (see Fig. 4 for $p = 3$ and [13]):

$$\begin{aligned}\mathbf{pq2_1} &= \{\mathbf{a},\mathbf{b},\mathbf{s} : \mathbf{a}^p = \mathbf{b}^q = \mathbf{asa}^{-1}\mathbf{s}^{-1} = \mathbf{babs}^{-1} = \mathbf{1}\} = \\ &= \{\mathbf{a},\mathbf{b} : \mathbf{a}^p = \mathbf{b}^q = \mathbf{ababa}^{-1}\mathbf{b}^{-1}\mathbf{a}^{-1}\mathbf{b}^{-1} = \mathbf{1}\}.\end{aligned} \tag{3.1}$$

Here $\mathbf{a}$ is a p-rotation about the fibre line through the origin (x axis), $\mathbf{b}$ is a q-rotation about the fibre line trough A_1 and $\mathbf{s} = \mathbf{bab}$ is a screw motion $\mathbf{s}: OA_1A_2 \to O^{\mathbf{s}}B_pB_1$. All these can be obtained by formulas (1.8) and (1.8'). Then we get the second presentation in (3.1), i.e. $\mathbf{abab} = \mathbf{baba} =: \tau$ is a fibre translation. Then $\mathbf{ab}$ is a $\mathbf{2_1}$ half-screw motion about $f_0 = HH^\tau$ (look at Fig. 4) that determines the fibre tarnslation τ above. This group in (3.1) surprisingly occurred in § 6 of our paper [7] at double links $K_{p,q}$. The coordinates of the vertices $A_1A_2A_3\dots A_p$ of the base figure and the corresponding vertices $B_1B_2B_3\dots B_p$ of the cover face can be computed for all given parameters by

$$\tanh(OA_1) = b := \sqrt{\frac{1 - \tan\frac{\pi}{p}\tan\frac{\pi}{q}}{1 + \tan\frac{\pi}{q}\tan\frac{\pi}{q}}}. \tag{3.2}$$

Moreover, the equation of the curve $c_{A_1A_2}$ can be determined as the foot points (see (1.4) and (1.9)) of the corresponding fibre lines. For example, the data of $\mathcal{P}_3(q)$ for some $6 < q \in \mathbb{N}$ are collected in Table 2 by Maple computations.

Table 2

(p,q)	b
$(3,7)$	≈ 0.30007426
$(3,8)$	≈ 0.40561640
$(3,9)$	≈ 0.47611091
$(3,10)$	≈ 0.50289355
$(3,50)$	≈ 0.89636657
$(3,1000)$	≈ 0.99457331
$(3,\infty)$	1

3.1 The volume of the bounded regular prisms $\mathcal{P}_p(q)$

The volume formula of a *sector-like* 3-dimensional domain $Vol(D(\Phi))$ can be computed by the metric tensor g_{ij} (1.13) in hyperboloid coordinates. This defined by the base figure D lying in the base plane (see Fig. 5) and by fibre translation τ given by (1.3) with parameter $\Phi \in \mathbb{R} \setminus \{0\}$.

Theorem 3.2 *Suppose we are given a sector-like region D illustrated in Fig. 5.a so a continuous function $r = r(\theta)$ where the radius r depends upon the angle θ. The volume of domain $D(\Phi))$ is derived by the following integral:*

$$\begin{gathered} Vol(D(\Phi)) = \int_D \frac{1}{2}\sinh(2r) dr\, d\theta\, d\phi = \\ = \int_0^{\Phi}\int_{\theta_1}^{\theta_2}\int_0^{r(\theta)} \frac{1}{2}\sinh(2r)\, dr\, d\theta\, d\phi = \Phi \int_{\theta_1}^{\theta_2} \frac{1}{4}(\cosh(2r(\theta)) - 1)\, d\theta. \end{gathered} \tag{3.3}$$

Let $\mathcal{T}_p(q)$ be a regular prism tiling above and let $\mathcal{P}_p(q)$ be one of its tiles. We get the following

Theorem 3.3 *The volume of a bounded regular prism $\mathcal{P}_p(q)$ $\Big(3 \le p \in \mathbb{N}, \frac{2p}{p-2} < q \in \mathbb{N}\Big)$ can be computed by the following formula:*

$$Vol(\mathcal{P}_p(q)) = Vol(D(p,q,\Phi)) \cdot p, \tag{3.4}$$

where $Vol(D(p,q,\Phi))$ *is the volume of the sector-like 3-dimensional domain that is given by the sector region* $OA_1A_2 \subset \mathcal{P}$ *(see Fig.5.b) and by* $\Phi = A_1B_1$*, the* $\widetilde{\mathbf{SL_2R}}$ *height of the prism, depending on* p, q*.*



a. b.

Figure 5: The sector region D

4 Ball packings to regular prism tilings

Sphere packing problems concern arrangements of non-overlapping equal spheres, rather balls, which fill a space. Space is the usual three-dimensional Euclidean space. However, ball (sphere) packing problems can be generalized to the other 3-dimensional Thurston geometries. But sometimes a difficult problem is – similarly to the hyperbolic space – the exact definition of the packing density. In [14] we extended the problem of finding the densest geodesic ball packing for the other 3-dimensional homogeneous geometries (Thurston geometries). In this paper we study the problem in $\widetilde{\mathbf{SL_2R}}$ geometry and develop a procedure for regular prism tilings and their above group $\mathbf{pq2_1}$ in (3.1).

We consider the above regular prism tilings $\mathcal{T}_p(q)$.

The *greatest ball* of centre K inside of $\mathcal{P}_p(q)$ is denoted by B_K, moreover let $\rho(K)$ denote the *radius* of B_K. The images of B_K in each tile of $\mathcal{T}_p(q)$ form a ball packing $\mathcal{B}_K$.

Definition 4.1 *The* density *of ball packing $\mathcal{B}_K$ is*

$$\delta(K) = \frac{Vol(B_K)}{Vol(\mathcal{P}_p(q))}.$$

Our problem is to find a point $K \in \mathcal{P}_p(q)$ and so the ball packing $\mathcal{B}_K$ where its density $\delta(K)$ is maximal. In this case the ball packing $\mathcal{B}_K$ is said to be *locally optimal*.

Our aim is to determine the maximal radius $\rho(K_{opt})$ of the balls, and the maximal density $\delta(\Gamma_p(q)) = \delta(K_{opt})$. The considered prism tiling does not have free parameters, thus

$$\delta(\Gamma_p(q)) = \max_K(\delta(K)). \tag{4.1}$$

4.1 The optimal ball packings to prism tilings $\mathcal{T}_p(q)$

Let $\mathcal{T}_p(q)$ be a regular prism tiling and let $\mathcal{P}_p(q)$ be one of its tiles which is given by its base figure $\mathcal{P}$ that is centered at the origin with vertices $A_1A_2A_3\dots A_p$ in the base plane of the model (see Fig. 7) and the corresponding vertices B_1B_2 $B_3\dots B_p$ and $C_1C_2C_3\dots C_p$ are generated by fibre translations τ and $-\tau$ given by (1.3) with parameter $\Phi \in \mathbb{R} \setminus \{0\}$, as above.

It is clear that the optimal ball B_K has to touch some faces of the corresponding prism $\mathcal{P}_p(q)$ and it is centered at the origin. The $Vol(\mathcal{P}_p(q))$ is given by the parameters p and q and $\Phi \geq \rho(K_{opt})$. The images of $\mathcal{P}_p(q)$ by our discrete group $\mathbf{pq2_1}$ covers the $\widetilde{\mathbf{SL_2R}}$ space without overlap. For the density of the packing it is sufficient to relate the volume of the optimal ball to that of the solid $\mathcal{P}_p(q)$ (see Definition 4.1 and (4.1)).

The maximal radius $\rho(K_{opt})$ of the balls can be determined for all possible parameters as the distance ρ between the origin and $c_{A_1A_2}$ (see Fig. 6), related to the half height $\Phi = \frac{\pi}{2} - \frac{\pi}{q} - \frac{\pi}{p}$ of the prism, also determined by p and q (see [7] and [13]).

The above locally densest geodesic ball packings can be determined for all regular prism tilings $\mathcal{T}_p(q)$ (p, q as above). For example, in Table 3 we have summarized some results to tilings $\mathcal{T}_3(q)$ $(6 < q \in \mathbb{N})$.



a. b.

Figure 6: The optimal half prism $A_1A_2A_3B_1B_2B_3$ with optimal half sphere for parameters $p = 3$, $q = 7$ with the maximal radius $\rho(K_{opt}) = \Phi = \frac{\pi}{2} - \frac{\pi}{p} - \frac{\pi}{q} < \rho$.

Table 3

(p,q)	$\rho(K_{opt})$	$Vol(B_K)$	$Vol(\mathcal{P}_p(q))$	$\delta(K_{opt})$
$(3,7)$	0.074800	0.001756	0.016785	0.104635
$(3,8)$	0.130899	0.009449	0.051404	0.183816
$(3,9)$	0.174533	0.022497	0.091385	0.246177
$(3,10)$	0.209440	0.039048	0.131595	0.296731
$(3,50)$	0.272672	0.087043	0.636918	0.136663
$(3,1000)$	0.274648	0.088981	0.812627	0.109498

4.2 Multiply transitive optimal ball packings under $\mathbf{pq2_1}$

We study only one case of the multiply transitive ball packings where the fundamental domain of the $\widetilde{\mathbf{SL_2R}}$ space group $\mathbf{pq2_1}$ is not a prism. Let the fundamental domains be Dirichlet- Voronoi cells where their centers are images of $K = O$.

These locally densest geodesic ball packings can be determined for all possible fixed parameters p and q $\left(3 \le p \in \mathbb{N},\ \frac{2p}{p-2} < q \in \mathbb{N}\right)$. The optimal radius $\rho(K_{opt})$ $(K = O)$ is

$$\rho(K_{opt}) = \min\left[\operatorname{artanh}(OA_1),\ \Phi = \frac{\pi}{2} - \frac{\pi}{p} - \frac{\pi}{q},\ \frac{d(O, O^{\mathbf{ab}})}{2}\right], \tag{4.2}$$



Figure 7: The optimal cell and ball for parameters $p = 3$ and $q = 7$.

where $d(O, O^{\mathbf{ab}})$ is the distance between O and $O^{\mathbf{ab}}$ by Definition 2.1.

In Table 4 we have summarized some results and we noticed that in the case $\mathcal{T}_6(9)$ we have obtained a very dense ball packing whose density is larger than ≈ 0.85327609 the universal hyperbolic 3-dimensional density upper bound by Böröczky and Florian [1]. Hence we have as a corollary

Table 4

(p,q)	$\rho(K_{opt})$	$Vol(B_K)$	$Vol(\mathcal{P}_p(q))$	$\delta(K_{opt})$
$(3,11)$	0.237999	0.057543	0.169931	0.338626
$(3,12)$	0.261799	0.076892	0.205617	0.373960
$(3,13)$	0.279134	0.093489	0.238467	0.392044
$(3,14)$	0.287083	0.101857	0.268561	0.379271
$(3,50)$	0.350810	0.188371	0.636918	0.295754
$(3,1000)$	0.370822	0.223543	0.812627	0.275087
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$(6,8)$	0.654498	1.350812	0.640122	0.823604
$\mathbf{(6,9)}$	$\mathbf{0.692287}$	$\mathbf{1.624770}$	$\mathbf{1.877243}$	$\mathbf{0.858262}$

Theorem 4.1 *The optimal ball packing to the* $\widetilde{\mathbf{SL_2R}}$ *group* $\mathbf{pq2_1}$ *above for* $p =$

6, $q = 9$ *with ball centres in the* p*-gonal rotation axis, is denser than any ball packing, related to its Dirichlet-Voronoi partition, in the hyperbolic space* $\mathbf{H}^3$.

Remark 4.1 *In a forthcoming paper we will study the above locally optimal geodesic ball packing arrangements and their densities for all possible parameters.*

Our projective method gives us a way of investigating the Thurston geometries. This suits to study and solve similar problems. (see e.g. [6], [9], [12], [14]).

References

[1] Böröczky, K. – Florian, A. Über die dichteste Kugelpackung im hyperbolischen Raum, *Acta Math. Acad. Sci. Hungar.*, (1964) **15** , 237–245.

[2] Divjak, B. – Erjavec, Z. – Szabolcs, B. – Szilágyi, B. Geodesics and geodesic spheres in $\widetilde{\mathbf{SL}(2,\mathbf{R})}$ geometry. *Math. Commun.,* **14/2** (2009), 413–424.

[3] E. Molnár – Szilágyi, B. Translation curves and their spheres in homogeneous geometries. *Publicationes Math. Debrecen.,* **78/2** (2011), 327–346.

[4] Molnár, E. The projective interpretation of the eight 3-dimensional homogeneous geometries. *Beiträge zur Algebra und Geometrie (Contributions to Algebra and Geometry),* **38** (1997) No. 2, 261–288.

[5] Molnár, E. – Szirmai, J. Symmetries in the 8 homogeneous 3-geometries. *Symmetry: Culture and Science,* **21/1-3** (2010), 87-117.

[6] Molnár, E. – Szirmai, J. Classification of $\mathbf{Sol}$ lattices. *Geometriae Dedicata*, **161/1** (2012), 251-275, DOI: 10.1007/s10711-012-9705-5.

[7] Molnár, E. – Szirmai, J. – Vesnin, A. Projective metric realizations of cone-manifolds with singularities along 2-bridge knots and links. *Journal of Geometry,* **95** (2009), 91-133.

[8] Scott, P. The geometries of 3-manifolds. *Bull. London Math. Soc.*, **15** (1983) 401–487.

[9] Szirmai, J. The densest geodesic ball packing by a type of $\mathbf{Nil}$ lattices. *Beiträge zur Algebra und Geometrie (Contributions to Algebra and Geometry),* **48(2)** (2007) 383–398.

[10] Szirmai, J. Geodesic ball packing in $\mathbf{S}^2\times\mathbf{R}$ space for generalized Coxeter space groups. *Beiträge zur Algebra und Geometrie (Contributions to Algebra and Geometry),* **52(2)** (2011), 413–430.

[11] Szirmai, J. Geodesic ball packing in $\mathbf{H}^2 \times \mathbf{R}$ space for generalized Coxeter space groups. *Math. Commun.,* **17/1** (2012), 151-170.

[12] Szirmai, J. Lattice-like translation ball packings in $\mathbf{Nil}$ space. *Publ. Math. Debrecen,* **80/3-4** (2012), 427–440 (DOI: 10.5486/PMD.2012.5117).

[13] Szirmai, J. Regular prism tilings in $\widetilde{\mathbf{SL_2R}}$ space. *Submitted Manuscript* (2012).

[14] Szirmai, J. A candidate to the densest packing with equal balls in the Thurston geometries. *Submitted Manuscript* (2012).

[15] Thurston, W. P. (and LEVY, S. editor) *Three-Dimensional Geometry and Topology.* Princeton University Press, Princeton, New Jersey, Vol **1** (1997).